

# Recent Ostracoda from estuarine systems of Ghana, western Africa


Peter Frenzel<sup>1\*</sup>, Lailah Gifty Akita<sup>2</sup>, Olga Gildeeva<sup>1</sup>, Maria Schott<sup>1</sup> & Mauro Alivernini<sup>1</sup>



UNIVERSITY OF GHANA  
DEPARTMENT OF MARINE AND FISHERIES SCIENCES



<sup>1</sup> Institute of Geosciences, Friedrich Schiller University Jena / <sup>2</sup> Department of Marine and Fisheries Sciences, University of Ghana






1 Aglaocypris gambiensis GH39 6 Neomonoceratina lidoensis GH39 11 Perissocytheridea GH36 16 Eucythereura? sp. GH39  
2 Cyprididae gen. et sp. GH35 7 Neomonoceratina GH39 12 Eucythere curta GH39 17 Neocaudites sp. GH35  
3 Paracypris GH39 8 Pontocythere sp. GH13 13 Paracytheridea GH36 18 Hemicytherura cellulosa GH36  
4 Candonidae gen. sp. GH39 9 Cyprinotus sp. GH46 14 Loxoconcha GH39  
5 Pseudoconcha GH49 10 Cyprideis remanei GH39 15 Chrysocythere foveostriata GH39

Here we present a faunistic overview of ostracods from ten estuarine systems covering the entire coast of Ghana. In total 58 surface samples and eight short cores were taken for the first study on brackish water ostracods from Ghana. The only previously published papers on Recent ostracods from Ghana are two studies about freshwater taxa.

We found 48 ostracod species, three platycopid and 45 podocopid species belonging to 35 genera. The most diverse families are Trachyleberididae and Hemicytheridae, the most diverse genus is *Puriana* with four different species. The fauna closely resembles the ostracod fauna from adjacent Nigeria and, to some degree, those from Gambia and Senegal but are distinctively different from those of the North African or South African coasts.

Ostracod diversity tends to decrease with salinity, highest values are found in the outer estuaries with polyhaline to euhaline conditions. High abundance of the brackish water taxa *Gibbaborchella kuznetsovae* (Omatso 1970) and *Cyprideis remanei* Klie 1940 indicates estuarine waters strongly impacted by organic pollution.



Ostracod taxa of Keta Lagoon and Densu Estuary. (1) *Basslerites punctata*; (2-3) *Pseudoconcha* sp.; (4-5) *Cyprideis remanei*; (6) *Paradoxostoma* sp.; (7) *Cytherella hanaii*; (8) *Neomonoceratina* sp. 2; (9) *Pontocythere* sp.; (10-11) *Loxoconcha lacunensis*; (12) *Cytherella* sp.; (13) *Puriana* aff. *interrasilis*; (14) *Puriana* cf. *rugosa*; (15) *Neomonoceratina* sp.; (16) *Neonesidea* sp.; (17) *Neomonoceratina* sp.; (18-19) *Pseudoconcha hartmanni*; (20-21) *Neomonoceratina* cf. *ikoroduensis*; (22) *Soudanella africana*; (24) *Puriana trituberculata* (25-26) *Eucytheridea*, gen. et sp. inc.; (27) *Chrysocythere foveostriata*; (28) *Neomonoceratina* cf. *ikoroduensis*; (29) *Perissocytheridea* (*Kroemmelbeinella*) *perfidiosa*; (30) *Puriana mediocostata*; (31) *Hermanites* cf. *foveolata*; (32) *Chrysocythere foveostriata*; (33) *Cleocythereis* sp.; (34) *Neonesidea* sp.; (35) *Carinocythereis* sp.; (36) *Neocaudites* sp.; (37) *Mackenziella lagosensis*; (38) *Gibbaborchella kuznetsovae*; (39) *Paracytheridea* sp.; (40) *Neocaudites lindersae*.

<b>Class Ostracoda Latreille 1802</b>
Subclass Platycopoda Sars 1866
Order Platycopina Sars 1866
Superfamily Cytherelloidea Sars 1866
Family Cytherellidae Sars 1866
<i>Cytherella hanaii</i> Omatsola 1970
<i>Cytherella</i> sp.
<i>Cytherelloidea? lobitoensis</i> Hartmann 1974
Subclass Podocopa Sars 1866
Order Podocopida Sars 1866
Superfamily Benthidae Sars 1865
Family Benthidae Sars 1865
<i>Neonesidea</i> sp.
Superfamily Cypridoidae Baird 1845
Family Candonidae Kaufmann 1900
<i>Aglaocypris gambiensis</i> Witte 1993
<i>Paracypris</i> sp.
Family Cyprididae Baird 1845
<i>Cyprinotus</i> sp.
<i>Heterocypris</i> sp.
Superfamily Cytheroidea Baird 1850
Family Cushmanidae Puri 1974
<i>Pontocythere</i> sp.
Family Cytheridae Baird 1850
<i>Pajenborchellina</i> sp.
Family Cytheridae Sars 1925
<i>Cyprideis remanei</i> Klie, 1940
<i>Perissocytheridea ebutemetaensis</i> (Omatsola 1970)
<i>Perissocytheridea</i> ( <i>Kroemmelbeinella</i> ) <i>libidinosa</i> (Witte 1986)
<i>Perissocytheridea</i> ( <i>Kroemmelbeinella</i> ) <i>perfidiosa</i> Witte 1993
Family Cytheridae Müller 1894
<i>Cytheropteron</i> sp.
<i>Cytherura</i> sp.
<i>Gibbaborchella kuznetsovae</i> (Omatsola 1970)
<i>Hemicytherura</i> sp.
<i>Semicythere duracina</i> Witte 1993
Family Eucytheridae Puri 1954
<i>Eucythere curta</i> Ruggieri, 1975
<i>Eucytherides</i> sp.
Family Hemicytheridae Puri 1953
<i>Hermanites</i> cf. <i>foveolata</i> Omatsola 1972
<i>Hermanites</i> sp.
<i>Neocaudites lindersae</i> Witte 1993
<i>Neocaudites</i> aff. <i>lindersae</i> sensu Witte 1993
<i>Puriana mediocostata</i> Omatsola 1972
<i>Puriana</i> aff. <i>interrasilis</i> sensu Omatsola 1972
<i>Puriana</i> cf. <i>rugosa</i> Omatsola 1972
<i>Puriana trituberculata</i> Omatsola 1972
Family Leptocytheridae Hanai 1857
<i>Lepocythere</i> sp.
Family Loxoconchidae Sars 1925
<i>Loxoconcha lacunensis</i> Omatsola 1970
<i>Pseudococha hartmanni</i> (Omatsola 1970)
<i>Pseudococha</i> sp.
Family Paracytheridae Puri 1957
<i>Paracytheridea</i> sp. A sensu Witte 1993
<i>Paracytheridae</i> gen. et sp. inc.
Family Paradoxostomatidae Brady and Norman 1889
<i>Paradoxostoma</i> sp.
Family Schizocytheridae Howe 1961
<i>Neomonoceratina ikoroduensis</i> Omatsola 1970
<i>Neomonoceratina</i> sp. 1
<i>Neomonoceratina</i> sp. 2
Family Trachyleberididae Sylvester-Bradley 1948
<i>Acanthocythereis</i> sp.
<i>Basslerites</i> ( <i>Loxoconcha</i> ) <i>punctata</i> Omatsola 1970
<i>Chrysocythere boldii</i> Omatsola 1970
<i>Chrysocythere foveostriata</i> (Brady 1870)
<i>Cleocythereis</i> sp.
<i>Reymontia ijejuorum</i> Omatsola 1970
<i>Ruggiera nigeriana?</i> Omatsola 1970
<i>Soudanella africana</i> Omatsola 1972
Ostracoda incertae sedis
<i>Mackenziella lagosensis</i> Omatsola 1972